




| Treatment Overview



“Without the ability to express ourselves through sexuality, and to form and maintain intimate relationships, we cannot achieve health.”


Good Sexual Energy

Decreases

- Stress
- Risk of ulcers
- Depression
- Pain
- Heart disease
- The incidence of cancer
- Incidence of myocardial infarction by 50% with twice weekly relations

Increases


- Sense of well being
- Longevity
- Immune system function
- Joy
- Tissue health & bladder function
- Self-esteem & happiness
- Quality of sleep



Causes of Erectile Dysfunction



Depression, anxiety, stress,
hyperextension, sleeping disorders,
pornography addiction



High blood pressure, stroke, high cholesterol,
clogged blood vessels (Atherosclerosis),
chronic heart problems



Alcohol, smoking, obstructive
pulmonary disease



Kidney disease like
renal failure or renal insufficiency,
certain prescription medications



Sexual boredom, abdominal or prostate surgery,
Parkinson's disease, Peyronie's disease –
development of scar tissue inside the penis,
treatment for prostate cancer
or enlarged prostate, declining testosterone



Diabetes, obesity, multiple sclerosis,
cerebral or spinal cord injuries that affect the
pelvic area or spine, cycling for long period




Phases of Sexual Response Cycle & Associated Sexual Dysfunctions

PHASES	CHARACTERISTICS	DYSFUNCTION
Desire	Reflects person's motivation, drives and personality; characterized by sexual fantasies and desire to have sex	Hypoactive sexual desire disorder; sexual aversion disorder
Excitement	Subjective sense of sexual pleasure and accompanying physiological responses (sexual flush, erection by vasocongestion, tightening and lifting of scrotal sac, increase size of testes)	Male erectile disorder; dyspareunia
Orgasm	Peaking of sexual pleasure, release of sexual tension and rhythmic contraction of perineal muscles and pelvic reproductive organs	Orgasmic disorder; premature ejaculation
Resolution	A sense of general relaxation, well-being and muscle relation	Post coital dysphoria; Post coital headache

Age & Erectile Dysfunction

Age Is a Major Risk Factor for Erectile Dysfunction



N=1209 noninstitutionalized men aged 40-70 years.
Feldman HA, et al.

Prevalence of Male Sexual Dysfunction

The Risk of Sexual Dysfunction increases with hypertension, diabetes, dyslipidemia, cardiovascular disease & smoking



Of men suffer from some form of ED




Of those who report ED are 40-60 years old



Of ED goes unreported because both practitioners and patients are not comfortable discussing their sexual health

Male Testosterone is needed for blood vessels to function



■ Testosterone Level ■ Testosterone Deficiency

Testosterone levels decline with age

Types of Pharmaceuticals (PDE-5 Inhibitors)

Medication	Viagra (sildenafil)	Levitra (vardenafil)	Cialis (tadalafil)	Stendra (avanafil)
Type	PDE-5 Inhibitor	PDE-5 Inhibitor	PDE-5 Inhibitor	PDE-5 Inhibitor
Dose	25-100mg	5-20mg	5-20mg	50-200mg
Peak Time	1 hour	42-54- minutes	2 hours	15-30 minutes
Gone from body	8-12 hours	8-12 hours	36 hours	8-12 hours
Contra-indicated	Nitrates	Nitrates	Nitrates	Nitrates
FDA Approval	3/29/98	8/20/03	2/02/04	4/01/12
Effects of eating and drinking	No food or drink 1-2 hours before	Not effected by food or alcohol	No food or drink 1-2 hours before	Not effected by food or alcohol
Side Effects	Headache, flushing, nasal congestion, abnormal vision, hearthburn, bloodshot eyes	Headache, flushing, nasal congestion, abnormal vision, hearthburn, bloodshot eyes	Headache, flushing, nasal congestion, abnormal vision, hearthburn, bloodshot eyes, backache, leg cramps	Headache, flushing, nasal congestion, abnormal vision, hearthburn, bloodshot eyes, backache, leg cramps
Other	For many patients the most potent. Negatives are shorter half-life (than Cialis), less effective when taken with fatty meal. Most side effects.	Greater selectivity (and thus usually fellow side effects) than Viagra. Slower to take effect when taken with fatty meal.	Greater selectivity (and thus usually fewer side effects) than Viagra.	Greater selectivity (and thus usually fellow side effects) than Viagra. Slower to take effect when taken with fatty meal.

Traditional Treatments (PDE-5 Inhibitors)



Pharmaceuticals Are Not For Everyone

Patients with the conditions listed below should avoid PDE-5 inhibitors

- Severe forms of heart failure
- Use of nitroglycerine
- Severe forms of heart disease, such as arrhythmias, unstable angina, a recent heart attack
- Very low blood pressure (hypotension)
- Uncontrolled high blood pressure (hypertension)
- Uncontrolled diabetes
- Recent history of stroke
- Retinitis pigmentosa (PDE-5 non-responders)



Side Effects of Pharmaceuticals



- Doesn't allow for spontaneity of intimacy
- Headache, flushing, runny nose, indigestion
- Back pain (Cialis) and indigestion
- Some men (<2%) experience prolonged or painful erections
- Visual problems (blurred vision, increased sensitivity to light, bluish haze) may occur


Clinical Studies

- Non-invasive treatment using Low-Intensity Extracorporeal Shockwave Therapy (LI-ESWT)
- Based on series of over 40 clinical studies showing 75+% success rate in treating ED



GAINSWave® is Safe, Proven & Effective

- Utilizes high frequency, low intensity soundwaves to increase blood flow to the penis
- Activates dormant cells, repairs aged blood vessels, stimulates growth of new blood vessels, removes decades of micro-plaque and activates new nerve tissue
- Uses a specific protocol designed to optimize efficacy, safety and results
- Drug- and surgery-free treatment performed in office
- Erections are firmer and overall sexual function is improved




What To Expect?

- Apply BLT numbing cream 30 minutes prior to your treatment
- Allow 20-30 minutes for completion of the procedure
- Immediate resumption of normal daily activities
- 75% of men experience positive results after 6 treatments


Additional Option - Platelet-Rich Plasma Shot (PRP Shot)

- After GAINSWave®, the PRP Shot is very effective to aid in healing the "trauma" site with growth factors
- PRP Shot incorporates harvesting & injection of one's own PRP into penis
- It is a natural process that the body does not reject, and is very successful



Additional Option-Penis Pump

- Use at home after GAINSWave® treatment for "workout" to increase blood flow
- Uses vacuum suction to pull blood into the penis to achieve natural, sustainable erections
- Over 90% of patients find relief with Vacuum Erection Device (VED), regardless of underlying cause of ED



GAINSWave®

The logo graphic consists of three wavy lines. The top line is blue, the middle line is yellow, and the bottom line is blue. They are arranged in a way that suggests a wave or a stylized 'W' shape.

GAINSWave.com

